

LAMB CREEP 16

MEDICATED

FOR THE DEVELOPMENT OF WEANED LAMBS

For the prevention of coccidiosis in young sheep caused by *Eimeria bakuensis*, *Eimeria crandallis*, *Eimeria ovinoidalis* and *Eimeria parva*.

ACTIVE DRUG INGREDIENT

Decoquinatate 27.2 g/ton

GUARANTEED ANALYSIS

Crude Protein, not less than	16.00%
(This includes not more than 1.0% equivalent protein from non-protein nitrogen)	
Crude Fat, not less than	4.00%
Crude Fiber, not more than	12.00%
Calcium, not less than	1.00%
Calcium, not more than	1.30%
Phosphorus, not less than	0.50%
Salt, not less than	0.50%
Salt, not more than	0.90%
Selenium, not less than	0.30 PPM
Vitamin A, not less than	6,000 I.U./lb.
Vitamin E, not less than	20 I.U./lb.

INGREDIENTS: Processed grain by-products, roughage products 12%, grain products, plant protein products, cane molasses, calcium carbonate, ammonium chloride 15lbs per ton, salt, vitamin A supplement, vitamin D3 supplement, vitamin E supplement, d-biotin, vitamin B12 supplement, choline chloride, riboflavin supplement, niacin supplement, d-calcium pantothenate, menadione sodium bisulfite complex, folic acid, L-ascorbyl-2-polyphosphate(vitamin C supplement), thiamine mononitrate, pyridoxine hydrochloride, manganese sulfate, zinc sulfate, ethylenediamine dihydriodide, cobalt sulfate, selenomethionine, and sodium selenite.

FEEDING DIRECTIONS: Feed **LAMB CREEP 16** to lambs for early growth and development. Feed continuously to provide decoquinatate at a daily dose of 22.7 mg./100 lbs. (0.5 mg./kg.) of bodyweight and feed for at least 28 days during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard.

<u>Bodyweight</u> <u>in pounds</u>	<u>*Daily feeding rate</u> <u>per head</u>
30	.5 lbs.
60	1.0 lbs.

*Feeding rate provides decoquinatate at a daily dose of 22.7 mg./100 lbs. (0.5 mg./kg.).

WARNING: DO NOT FEED TO SHEEP PRODUCING MILK FOR FOOD.

Manufactured by
LYSSY & ECKEL, INC.
POTH, TEXAS 78147

Net Wt. 50 lbs. (22.6 kg.)

842