

2 : 1 "LP" HI-MAG MINERAL

MEDICATED

(For Ruminants Only)

FOR STOCKER CATTLE ON PASTURE

ACTIVE DRUG INGREDIENT

Lasalocid 1428 gr/ton

Pasture Cattle (Slaughter, Stocker, Feeder Cattle, and Dairy and Beef Replacement Heifers) For increased rate of weight gain and for control of coccidiosis caused by *Eimeria bovis* and *E. zuernii*.

NOTE: Coccidiosis may occur when young pasture cattle are co-mingled with adult cattle passing coccidial oocysts.

CAUTION: Do not allow horses or other equines access to premixes or supplements containing lasalocid, as ingestion may be fatal.

The safety of lasalocid in unapproved species has not been established.

Feeding undiluted or mixing errors resulting in excessive concentrations of lasalocid could be fatal to cattle and sheep.

When mixing and handling lasalocid premix, use protective clothing, impervious gloves and a dust mask. Avoid contact with eyes. Operators should wash thoroughly with soap and water after handling.

WARNING: A withdrawal period has not been established for this product in preruminating cattle. Do not use in calves to be processed for veal.

GUARANTEED ANALYSIS

Calcium, not less than	15.00%
Calcium, not more than	1.00%
Phosphorus, not less than	7.50%
Salt, not less than	14.00%
Salt, not more than	16.00%
Potassium, not less than	1.85%
Magnesium, not less than	2.50%
Zinc, not less than	4500 ppm
Copper, not less than	1500 ppm
Selenium, not less than	30 ppm
Vitamin A, not less than	100,000 I.U./lb.
Vitamin D, not less than	40,000 I.U./lb.

INGREDIENTS: Dicalcium phosphate, monocalcium phosphate, calcium carbonate, salt, plant protein products, potassium sulfate, magnesium sulfate, sulfur, magnesium oxide, potassium chloride, cane molasses, iron oxide, manganese sulfate, zinc sulfate, copper sulfate, ethylenediamine dihydriodide, cobalt sulfate, sodium selenite, vitamin A supplement, vitamin D3 supplement, vitamin E supplement and artificial flavor.

FEEDING DIRECTIONS:

FOR INCREASED RATE OF WEIGHT GAIN: Feed 1.4 oz. to 4.5 oz to provide not less than 60 mg. nor more than 200 mg. lasalocid per head per day.

FOR CONTROL OF COCCIDIOSIS: Hand feed continuously at the rate of 1 mg. lasalocid per 2.2 pounds of body weight per day in cattle up to 800 pounds (maximum 360 mg. lasalocid per day).

Manufactured by
LYSSY & ECKEL, INC.
POTH, TEXAS 78147
Net wt. 50 lbs. (22.6 kg.)