

COCCI-BAN

MEDICATED

As an aid in treatment of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii*.

ACTIVE DRUG INGREDIENT

Amprolium 445mg./lb.

GUARANTEED ANALYSIS

Crude Protein, not less than	12.00%
Crude Fat, not less than	3.00%
Crude Fiber, not more than	14.00%
Calcium, not less than	0.10%
Calcium, not more than	0.30%
Phosphorous, not less than	0.40%
Salt	none added
Potassium, not less than	0.70%
Vitamin A	none added

INGREDIENTS: Grain products, roughage products 15%, processed grain by-products, plant protein products, cane molasses, autolyzed brewers dried yeast, extracted citric acid press cake, yucca schidegera, active dry yeast, brewers dried yeast, brewers dried grains, saccharomyces cerevisiae fermentation extract, dried aspergillus oryzae fermentation extract, dried bacillus licheniformis fermentation extract, dried bacillus subtilus fermentation extract and artificial flavors

FEEDING DIRECTIONS: Feed **COCCI-BAN** to beef cattle diagnosed with coccidiosis to provide 10mg./kg. bodyweight for 5 days.

- Feed **COCCI-BAN** at the first clinical signs of coccidiosis, such as diarrhea and dehydration.

CATTLE WEIGHT	COCCI-BAN HEAD/DAY
500#	5
600#	6

WARNING: Withdraw 24 hours before slaughter. Use as sole source of Amprolium. A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.

**Manufactured by
LYSSY & ECKEL, INC.
POTH, TEXAS 78147
Net Wt. 50 lbs. (22.6 kg)**