

PASTURE GOAT

16

MEDICATED

FOR FREE-RANGING GOATS ON PASTURE

For the prevention of coccidiosis in young goats caused by *E. christenseni* and *E. ninakohlyakimovae*.

ACTIVE DRUG INGREDIENT

Decoquinatate 27.2 g/ton

GUARANTEED ANALYSIS

Crude Protein, not less than	16.00%
Crude Fat, not less than	4.00%
Crude Fiber, not more than	12.00%
Calcium, not less than	0.80%
Calcium, not more than	1.20%
Phosphorus, not less than	0.50%
Salt, not less than	0.80%
Salt, not more than	1.00%
Potassium, not less than	1.00%
Copper, not less than	60 ppm
Selenium, not less than	0.60 ppm
Vitamin A, not less than	20,000 I.U./lb.
Vitamin E, not less than	20 I.U./lb.

INGREDIENTS: Processed grain by-products, grain products, plant protein products, forage products, cane molasses, calcium carbonate, ammonium chloride, salt, dicalcium phosphate, monocalcium phosphate, vitamin A supplement, vitamin D3 supplement, vitamin E supplement, vitamin B12 complex, riboflavin supplement, niacin supplement, d-calcium pantothenate, menadione sodium bisulfite complex, folic acid, thiamine mononitrate, d-biotin, manganese sulfate, manganese amino acid complex, zinc sulfate, zinc amino acid complex, copper sulfate, copper amino acid complex, ethylenediamine dihydriodide, cobalt sulfate, cobalt glucoheptonate, selenomethionine, and sodium selenite.

FEEDING DIRECTIONS: Feed **PASTURE GOAT 16** to supplement free-ranging goats on native pasture. Feed continuously to provide decoquinatate at a daily dose of 22.7 mg./100 lbs.(0.5 mg./kg.) of bodyweight and feed for at least 28 days during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard. Adequate forage and water should be maintained at all times.

Goat Bodyweight in pounds

30
60
90
120

Daily Feeding Rate per head

0.50 lb.
1.00 lb.
1.50 lb.
2.00 lb.

Manufactured by
LYSSY & ECKEL, INC.
POTH, TEXAS 78147
Net Wt. 50 lbs. (22.6 kg.)