

SHOW GOAT

16/5

MEDICATED

FOR GOATS BEING FITTED FOR SHOW

For the prevention of coccidiosis in young goats caused by *Eimeria christensenii* and *Eimeria ninakohlyakimovae*.

ACTIVE DRUG INGREDIENT

Decoquinatate 27.2 g/ton

GUARANTEED ANALYSIS

Crude Protein, not less than	16.00%
(This includes not more than 1.0% equivalent protein from ammonium chloride)	
Crude Fat, not less than	5.00%
Crude Fiber, not more than	14.00%
Calcium, not less than	0.70%
Calcium, not more than	1.10%
Phosphorus, not less than	0.50%
Potassium, not less than	1.00%
Salt, not less than	0.50%
Salt, not more than	0.90%
Copper, not less than	20 PPM
Selenium, not less than	0.30 PPM
Vitamin A, not less than	20,000 I.U./lb.
Vitamin D, not less than	4,000 I.U./lb.
Vitamin E, not less than	50 I.U./lb.

INGREDIENTS: Ground corn, stabilized rice bran, alfalfa leaf meal, ground barley, soybean meal, cottonseed hulls 9%, beet pulp, flaxseed, cane molasses, calcium carbonate, menhaden fish meal, dried saccharomyces cerevisiae fermentation product, blood meal, ammonium chloride, dicalcium phosphate, monocalcium phosphate, dried marine algae, salt, potassium magnesium sulfate, yucca schidegera, magnesium oxide, vitamin A supplement, vitamin D3 supplement, vitamin E supplement, hemicellulose extract, carvacrol, autolyzed brewers dried yeast, extracted citric acid presscake, active dry yeast, brewers dried yeast, saccharomyces cerevisiae fermentation extract, dried aspergillus oryzae fermentation extract, dried bacillus licheniformis fermentation extract, dried bacillus subtilis fermentation extract, d-biotin, vitamin B12 supplement, riboflavin supplement, niacin supplement, d-calcium pantothenate, choline chloride, menadione sodium bisulfite complex, folic acid, thiamine mononitrate, pyridoxine hydrochloride, manganese sulfate, manganese amino acid complex, zinc sulfate, zinc amino acid complex, ethylenediamine dihydriodide, cobalt sulfate, cobalt glucoheptonate, copper sulfate, copper amino acid complex, sodium selenite, selenium yeast and artificial flavors.

FEEDING DIRECTIONS: Feed **SHOW GOAT 16/5** to goats being fitted for show. Feed continuously to provide decoquinatate at a daily dose of 22.7 mg./100 lbs. (0.5 mg./kg.) of bodyweight and feed for at least 28 days during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard.

<u>Goat Bodyweight</u> <u>in pounds</u>	<u>*Daily feeding rate</u> <u>per head</u>
30	1.0
60	2.0
90	3.0
120	4.0

*Feeding rate provides decoquinatate at a daily dose of 22.7 mg./100 lbs. (0.5 mg./kg.).

WARNING: DO NOT FEED TO GOATS PRODUCING MILK FOR FOOD.

Manufactured by
LYSSY & ECKEL, INC.
POTH, TEXAS 78147
Net Wt. 50 lbs. (22.6 kg.)