



# BREED BACK

## 20



### FOR COWS, HEIFERS AND BULLS WARMING UP FOR BREEDING

#### GUARANTEED ANALYSIS

Crude Protein, min .....	20.00%	Zinc (Z), min .....	1850 ppm
Crude Fat, min .....	5.00%	Manganese (Mn), min .....	1500 ppm
Crude Fiber, max .....	2.00%	Copper (Cu), min .....	.650 ppm
Calcium (Ca), min .....	2.00%	Cobalt (Co), min .....	40 ppm
Calcium (Ca), max .....	3.00%	Iodine (I), min .....	40 ppm
Phosphorus (P), min .....	1.50%	Selenium (Se), min .....	5 ppm
Salt (NaCl) .....	0 added	Vitamin A, min .....	100,000 IU/lb
Magnesium (Mg), min .....	2.00%	Vitamin D, min .....	10,000 IU/lb
Potassium (K), min .....	3.00%	Vitamin E, min .....	150 IU/lb
		Total Sugars (as invert), min .....	29.00%

#### INGREDIENTS

Cane Molasses, Hydrolyzed Feather Meal, Cottonseed Meal, Monocalcium Phosphate, Vegetable Oil, Magnesium Oxide, Calcium Carbonate, Manganese Amino Acid Complex, Zinc Amino Acid Complex, Copper Amino Acid Complex, Zinc Sulfate, Manganese Sulfate, Cobalt Glucoheptonate, Copper Sulfate, Sodium Selenite, Vitamin E Supplement, Vitamin A Supplement, Vitamin D3 Supplement and Ethylenediamine Dihydriodide

Manufactured By:  
Hudson Livestock Supplements, Inc.  
8150 Thompson Rd  
Miles, TX 76861  
(325) 659-3992 \* 1-800-750-9608

#### FEEDING RECOMMENDATIONS

Consumption will vary depending upon the condition of the animals, quality and quantity of forages being fed, weather conditions, and most importantly the feeding location of the tubs with respect to loafing, grazing, feeding, and watering areas.

Cattle will consume approximately 1/2 to 1 pound per head daily based upon a 1,000 lb. animal. Tubs should be distributed at the rate of one tub per 20-25 head. Place the tubs where animals congregate (loafing, grazing, feeding, and watering areas). Feed the tubs continuously along with plentiful sources of forage and clean, fresh water. Provide free-choice salt at all times. When feedstuffs are limited, additional nutrients may be required. Contact your feed representative for specific recommendations.

**INTRODUCTION PERIOD:** During the first week of use, place the tubs approximately 100 feet from loafing and/or watering areas. As livestock become acclimated to the presence of the tubs, repositioning may be necessary to achieve proper consumption. Move tubs closer to increase, further away to decrease consumption. If overconsumption occurs for more than two weeks and repositioning does not correct the situation, remove tubs and evaluate the overall feeding program.

#### CAUTION:

**Use as directed. This product contains high levels of copper. Do not feed to sheep or any species that have a low tolerance to supplemental copper.**

NET WEIGHT 225 lb (102.06 kg)