

# SHOW CALF STRESS REGULATOR 10



# FOR REDUCING STRESS IN SHOW CALVES EXCELLING IN HIGH PERFORMANCE ENVIRONMENTS

### GUARANTEED ANALYSIS

Crude Protein, min	Cobalt (Co), min
Crude Fat, min	Iodine (I), min
Crude Fiber, max	Selenium (Se), min
Calcium (Ca), min	Vitamin A, min
Calcium (Ca), max	Vitamin D, min
Phosphorus (P), min	Vitamin E, min
Magnesium (Mg), min	Vitamin B12, min5 mgc/lb
Potassium (K), min	Choline, min
Zinc (Z), min	Riboflavin, min
Manganese (Mn), min 1200 ppm	Niacin, min
Copper (Cu), min 400 ppm	Total Sugars (as invert), min

# INGREDIENTS

Cane Molasses, Soybean Meal, Cottonseed Meal, Calcium Carbonate, Vegetable Oil, Sea Kelp, Autolyzed Brewers Dried Yeast, Extracted Citric Acid Presscake, Active Dry Yeast, Brewers Dried Yeast, Saccharomyces Cerevisiae Fermentation Extract, Dried Aspergillus Oryzae Fermentation Extract, Dried Bacillus Licheniformis Fermentation Extract, Dried Bacillus Subtillis Fermentation Extract, Saccharomyces Cerevisiae Yeast, Monocalcium Phosphate, Magnesium Oxide, Manganese Amino Acid Complex, Zinc Amino Acid Complex, Magnesium Proteinate, Eugenol, Cinnamaldehyde, Anethole, Capsicum, Vitamin E Supplement, Precision Niacin, Choline Chloride, Vitamin B12 Supplement, Calcium Pantothenate, Riboflavin Supplement, Vitamin A Supplement, Menadione Sodium Bisulfite Complex, Vitamin D3 Supplement, Thiamine Mononitrate, Biotin, Ascorbic Acid, Pyridoxine Hydrochloride, Folic Acid, Copper Amino Acid Complex, Zinc Sulfate, Cobalt Glucoheptonate, Selenium Yeast, Sodium Selenite, Manganese Sulfate and Copper Sulfate

### FEEDING RECOMMENDATIONS

Consumption will vary depending upon the condition of the animals, quality and quantity of forages being fed, weather conditions, and most importantly the feeding location of the tubs with respect to loafing, grazing, feeding, and watering areas.

Cattle will consume approximately 1/2 to 1 pound per head daily based upon a 1,000 lb. animal. Tubs should be distributed at the rate of one tub per 2-5 head. Place the tubs where animals congregate (loafing, grazing, feeding, and watering areas). Feed the tubs continuously along with plentiful sources of forage and clean, fresh water. Provide free-choice salt at all times. When feedstuffs are limited, additional nutrients may be required. Contact your feed representative for specific recommendations.

**INTRODUCTION PERIOD:** During the first week of use, place the tubs approximately 100 feet from loafing and/ or watering areas. As livestock become acclimated to the presence of the tubs, repositioning may be necessary to achieve proper consumption. Move tubs closer to increase, further away to decrease consumption. If overconsumption occurs for more than two weeks and repositioning does not correct the situation, remove tubs and evaluate the overall feeding program.

# CAUTION:

Use as directed. This product contains high levels of copper. Do not feed to sheep or any species that have a low tolerance to supplemental copper.

