

ZOO & WILDLIFE WORMER 20X10 2X

FOR RUMINANTS RAISED IN CONFINEMENT
(Subfamilies antilopinae, hippotraginae and caprinae)

For the removal and control of: Small stomach worm (*Trichostrongylus spp.*), thread necked intestinal worm (*Nematodirus spp.*), barberpole worm (*Haemonchus spp.*), and whipworm (*Trichuris spp.*).

LIMITATIONS: Feed as per use level chart over a 3-day period. Retreatment may be needed after 6 weeks. Do not use 14 days before or during hunting season.

NOTE: Consult veterinarian for assistance in the diagnosis, treatment and control of parasitism.

ACTIVE DRUG INGREDIENT

Fenbendazole 45.0 mg./lb.

GUARANTEED ANALYSIS

Crude Protein, not less than 20.00%
 Crude Fat, not less than 4.00%
 Crude Fiber, not more than 12.00%
 Calcium, not less than 1.80%
 Calcium, not more than 2.20%
 Phosphorus, not less than 1.10%
 Salt, not less than 0.50%
 Salt, not more than 0.80%
 Zinc, not less than 600 PPM
 Manganese, not less than 400 PPM
 Copper, not less than 150 PPM
 Vitamin A, not less than 20,000 I.U./lb.
 Vitamin E, not less than 50 I.U./lb.

INGREDIENTS: Processed grain by-products, plant protein products, forage products, grain products, cane molasses, dicalcium phosphate, monocalcium phosphate, calcium carbonate, salt, vitamin A supplement, vitamin D3 supplement, vitamin E supplement, zinc sulfate, zinc amino acid complex, manganese sulfate, manganese amino acid complex, copper amino acid complex, ethylenediamine dihydride, copper sulfate, cobalt sulfate, sodium selenite, cobalt glucoheptonate and artificial flavor.

FEEDING DIRECTIONS: Feed at the rates recommended below to provide 2.5mg./kg. of bodyweight per day for 3 days.

BODYWEIGHT	ZOO & WILDLIFE 20 WORMER/HEAD/DAY FOR 3 DAYS
40	1.00
80	2.00
120	3.00
160	4.00
200	5.00

Manufactured By
LYSSY & ECKEL, INC.
POTH, TEXAS 78147

Net Wt. _____