

SAFEGUARD DE-WORMER PELLETS MEDICATED

FOR BEEF AND DAIRY CATTLE

For the removal and control of lungworms (*Dictyocaulus viviparus*); barberpole worms (*Haemonchus contortus*); brown stomach worms (*Ostertagia ostertagi*); small stomach worms (*Trichostrongylus axei*); hookworms (*Bunostomum phlebotomum*); thread-necked intestinal worms (*Nematodirus helvetianus*); small intestinal worms (*Cooperia punctata* and *Cooperia oncophora*); bankrupt worms (*Trichostrongylus colubriformis*); and nodular worms (*Oesophagostomum radiatum*).

LIMITATIONS: In free-choice cattle feed, feed a total of 2.27 mg. Fenbendazole per pound of bodyweight to cattle over a 3-6 day period. Retreatment may be needed after 4-6 weeks. Cattle must not be slaughtered within 13 days following last treatment. For dairy cattle the milk discard time is zero hours. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. Note: Consult veterinarian for assistance in the diagnosis, treatment and control of parasitism.

ACTIVE DRUG INGREDIENT

Fenbendazole. 227.0 mg./lb.

GUARANTEED ANALYSIS

Crude Protein, not less than	14.00%
Crude Fat, not less than.	3.00%
Crude Fiber, not more than	16.00%
Calcium, not less than	0.80%
Calcium, not more than	1.20%
Phosphorus, not less than	0.60%
Salt, not less than.	0.50%
Salt, not more than.	0.80%

INGREDIENTS: Processed grain by-products, forage products, plant protein products, cane molasses, calcium carbonate, yeast culture, salt, cinnamaldehyde, anethole, eugenol, carvacrol, capsaicin and artificial flavors.

FEEDING DIRECTIONS: Feed **SAFEGUARD DE-WORMER PELLETS** at the rates recommended below to provide 2.27 mg./lb. bodyweight.

Bodyweight	SAFEGUARD DE-WORMER PELLETS
100	1.00
200	2.00
300	3.00
400	4.00
500	5.00
600	6.00
700	7.00
800	8.00

**Manufactured by
LYSSY & ECKEL, INC.
POTH, TEXAS 78147
Net Wt. 50 lbs. (22.6 kg.)**