

ROCKY TOP 14:7 + ALTOSID

A nutritional supplement containing S-Methoprene* Insect Growth Regulator for continuous feeding to cattle through the horn fly season to prevent adult horn fly emergence from manure of treated cattle.

ACTIVE INGREDIENT:

(S)- Methoprene (CAS #65733-16-6)..... 0.014%

OTHER INGREDIENTS:..... **99.986%**

TOTAL..... 100.00%

*As the Insect Growth Regulator ALTOSID®

EPA Reg. No. 89459-88-104656

EPA Est. No. 104656-TX-1

GUARANTEED ANALYSIS

Calcium, not less than	12.00%
Calcium, not more than	16.00%
Phosphorus, not less than	7.00%
Salt, not less than	16.00%
Salt, not more than	19.00%
Potassium, not less than	1.00%
Magnesium, not less than	2.00%
Zinc, not less than	4,750 PPM
Manganese, not less than	3,000 PPM
Copper, not less than	2,000 PPM
Selenium, not less than	25 PPM
Iodine, not less than.....	100 PPM
Vitamin A, not less than.....	375,000 I.U./lb.
Vitamin E, not less than	1,600 I.U./lb.
Thiamine, not less than	850 mg./lb.

FEED INGREDIENTS

Dicalcium phosphate, monocalcium phosphate, calcium carbonate, salt, yeast culture, magnesium oxide, cane molasses, potassium chloride, mineral oil, vitamin A supplement, vitamin D3 supplement, vitamin E supplement, thiamine mononitrate, eugenol, capsicum oleoresin, cinnamaldehyde, zinc hydroxychloride, zinc amino acid complex, manganese hydroxychloride, manganese amino acid complex, basic copper chloride, copper amino acid complex, sodium selenite, selenium yeast, cobalt glucoheptonate, ethylenediamine dihydriodide and artificial flavors.

**KEEP OUT OF REACH OF CHILDREN
WARNING**

See additional Precautionary Statements

PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING**

Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

FIRST AID:	
If in Eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact the National Poison Control Center at [1-800-222-1222], 24 hrs./day, 7 days/wk. for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION

This product is to be used as a cattle feed-through pesticide for the control of horn flies. All feed ingredients listed on this product label must adhere to the specifications and limits (if applicable) ascribed by the Food and Drug Administration (FDA) for the use of pesticides in cattle feed.

This product contains ethoxyquin, a preservative. The amount of ethoxyquin may vary according to the individual manufacturer's formulation and will not exceed the 150-ppm maximum limit established for ethoxyquin by FDA, per 21 CFR 573.380.

FEEDING AND MANAGEMENT INSTRUCTIONS:

Feed this product to cattle only. When used as directed, **ROCKY TOP 14:7 + ALTOSID** will provide sufficient (S)-Methoprene insect growth regulator to prevent the emergence of adult horn flies from manure of treated cattle. Existing adult horn flies will not be affected. Start feeding before horn flies appear and continue use until cold weather marks the end of the horn fly season. Allow free choice to cattle. Cattle should consume an average of 0.286 ounces of **ROCKY TOP 14:7 + ALTOSID** per 100 lbs. of cattle body weight per day, which is equivalent to 4 ounces per day for a 1,400-pound cow.

1. Assure cattle are not starved for nutrients in **ROCKY TOP 14:7 + ALTOSID** by providing them prior to feeding **ROCKY TOP 14:7 + ALTOSID**.
2. This product contains added copper. Do not feed to cattle in close proximity to sheep or other species that have a low tolerance to supplemental copper.
3. Offer cattle 50 pounds of this product for every 15 head of cattle.
4. Remove all other self-limiting nutritional supplements while cattle are consuming this product.
5. Locate **ROCKY TOP 14:7 + ALTOSID** where cattle congregate.
6. Replenish **ROCKY TOP 14:7 + ALTOSID** on a regular basis as needed.
7. If intake of **ROCKY TOP 14:7 + ALTOSID** is below 0.286 ounces per 100 pounds cattle body weight per day, increase the number of feeding locations to areas more frequented by cattle.
8. If intake of **ROCKY TOP 14:7 + ALTOSID** is above 0.286 ounces per 100 pounds cattle body weight per day, reduce the number of feeding locations and/or change locations to water sources less frequented by cattle.
9. Introduction of this product after adult horn fly infestation is established will require treatment of cattle with adulticides if elimination of the adult fly population is desired.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store this product in a cool dry place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Completely empty bag, then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

ALTOSID® Insect Growth Regulator is a registered trademark of Wellmark International.

**Manufactured by
LYSSY & ECKEL, INC.
POTH, TX 78147**

33089

Net Wt. 50 lbs. (22.6 kg.)