



WEAN CALF STRESS 12/4



FOR BEEF CATTLE

GUARANTEED ANALYSIS

Crude Protein, min	12.00%	Zinc (Z), min.....	2400 ppm
Crude Fat, min.....	4.00%	Manganese (Mn), min	1300 ppm
Crude Fiber, max.....	3.50%	Copper (Cu), min	750 ppm
Calcium (Ca), min	2.00%	Cobalt (Co), min	60 ppm
Calcium (Ca), max	3.00%	Iodine (I), min	40 ppm
Phosphorus (P), min	1.00%	Selenium (Se), min.....	13 ppm
Salt (NaCl), min	1.50%	Vitamin A, min	200,000 IU/lb
Salt (NaCl), max.....	2.50%	Vitamin D, min	20,000 IU/lb
Magnesium (Mg), min.....	3.00%	Vitamin E, min.....	800 IU/lb
Potassium (K), min.....	2.80%	Total Sugars (as invert), min	25.00%

INGREDIENTS

Cane Molasses, Cottonseed Meal, Saccharomyces Cerevisiae Fermentation Extract, Magnesium Oxide, Monocalcium Phosphate, Vegetable Oil, Salt, Calcium Carbonate, Zinc Amino Acid Complex, Propionic Acid (preservative), Cobalt Carbonate, Vitamin E Supplement, Manganese Hydroxychloride, Selenium Yeast, Sodium Selenite, Basic Copper Chloride, Vitamin A Supplement, Vitamin D3 Supplement and Ethylenediamine Dihydriodide

Manufactured By:
Hudson Livestock Supplements, Inc.
8150 Thompson Rd
Miles, TX 76861
(325) 659-3992 * 1-800-750-9608

FEEDING RECOMMENDATIONS

Consumption will vary depending upon the condition of the animals, quality and quantity of forages being fed, weather conditions, and most importantly the feeding location of the tubs with respect to loafing, grazing, feeding, and watering areas.

Cattle will consume approximately 1/2 to 1 pound per head daily based upon a 1,000 lb. animal. Tubs should be distributed at the rate of one tub per 20-25 head. Place the tubs where animals congregate (loafing, grazing, feeding, and watering areas). Feed the tubs continuously along with plentiful sources of forage and clean, fresh water. Provide free-choice salt at all times. When feedstuffs are limited, additional nutrients may be required. Contact your feed representative for specific recommendations.

INTRODUCTION PERIOD: During the first week of use, place the tubs approximately 100 feet from loafing and/or watering areas. As livestock become acclimated to the presence of the tubs, repositioning may be necessary to achieve proper consumption. Move tubs closer to increase, further away to decrease consumption. If overconsumption occurs for more than two weeks and repositioning does not correct the situation, remove tubs and evaluate the overall feeding program.

CAUTION:

Use as directed. This product contains high levels of copper. Do not feed to sheep or any species that have a low tolerance to supplemental copper.

NO SULFATES

NO OXIDES

NET WEIGHT 125 lb (56.70 kg)