



RUMEN FUEL BREED BACK 20



FOR BEEF CATTLE ON NATIVE AND IMPROVED PASTURE

GUARANTEED ANALYSIS

Crude Protein, min	20.00%	Potassium (K), min	2.50%
Not more than 6% equivalent protein from non-protein nitrogen		Zinc (Z), min	1800 ppm
Crude Fat, min.....	3.50%	Manganese (Mn), min	1600 ppm
Crude Fiber, max.....	1.25%	Copper (Cu), min	750 ppm
Calcium (Ca), min	2.00%	Cobalt (Co), min	10 ppm
Calcium (Ca), max	3.00%	Iodine (I), min	25 ppm
Phosphorus (P), min	2.00%	Selenium (Se), min.....	7.5 ppm
Salt (NaCl)	0 added	Vitamin A, min	150,000 IU/lb
Magnesium (Mg), min.....	2.00%	Vitamin D, min	20,000 IU/lb
		Vitamin E, min.....	400 IU/lb
		Total Sugars (as invert), min	25.00%

INGREDIENTS

Cane Molasses, Soybean Meal, Monocalcium Phosphate, Vegetable Oil, Magnesium Oxide, Urea, Zinc Amino Acid Complex, Manganese Hydroxychloride, Copper Amino Acid Complex, Vitamin E Supplement, Zinc Hydroxychloride, Sodium Selenite, Cobalt Carbonate, Selenium Yeast, Vitamin A Supplement, Vitamin D3 Supplement, Basic Copper Chloride and Ethylenediamine Dihydriodide

Manufactured By:
Hudson Livestock Supplements, Inc.
8150 Thompson Rd
Miles, TX 76861
(325) 659-3992 * 1-800-750-9608

FEEDING RECOMMENDATIONS

Consumption will vary depending upon the condition of the animals, quality and quantity of forages being fed, weather conditions, and most importantly the feeding location of the tubs with respect to loafing, grazing, feeding, and watering areas.

Cattle will consume approximately 1/2 to 1 pound per head daily based upon a 1,000 lb. animal. Tubs should be distributed at the rate of one tub per 20-25 head. Place the tubs where animals congregate (loafing, grazing, feeding, and watering areas). Feed the tubs continuously along with plentiful sources of forage and clean, fresh water. Provide free-choice salt at all times. When feedstuffs are limited, additional nutrients may be required. Contact your feed representative for specific recommendations.

INTRODUCTION PERIOD: During the first week of use, place the tubs approximately 100 feet from loafing and/or watering areas. As livestock become acclimated to the presence of the tubs, repositioning may be necessary to achieve proper consumption. Move tubs closer to increase, further away to decrease consumption. If overconsumption occurs for more than two weeks and repositioning does not correct the situation, remove tubs and evaluate the overall feeding program.

CAUTION:

Use as directed. For ruminants only. Do not feed to equine. Do not feed to sheep or any species that have a low tolerance to supplemental copper.

**NO SULFATES
NO OXIDES**



NET WEIGHT 225 lb (102.06 kg)