



STAMINA EQUINE 12



FOR GROWTH AND MAINTENANCE OF ALL EQUINE CLASSES

GUARANTEED ANALYSIS

Crude Protein, min	12.00%	Selenium (Se), min.....	3.1 ppm
Crude Fat, min.....	4.50%	Zinc (Z), min.....	1000 ppm
Crude Fiber, max.....	1.50%	Vitamin A, min	90,000 IU/lb
Calcium (Ca), min	3.00%	Vitamin D, min	20,000 IU/lb
Calcium (Ca), max	4.00%	Vitamin E, min	90 IU/lb
Phosphorus (P), min	2.00%	Menadione, min	30 mg/lb
Magnesium (Mg), min.....	3.00%	Vitamin B12, min.....	400 mcg/lb
Potassium (K), min.....	3.00%	Biotin, min	0.5 mg/lb
Sulphur (S), min	0.40%	Choline, min	800 mg/lb
Copper (Cu), min	100 ppm	Niacin, min	900 mg/lb
Cobalt (Co), min.....	27 ppm	d-Pantothenic Acid, min.....	250 mg/lb
Iron (Fe), min	500 ppm	Riboflavin, min	100 mg/lb
Iodine (I), min	13 ppm	Thiamine, min	30 mg/lb
Manganese (Mn), min	800 ppm	Ascorbic Acid, min	9 mg/lb
		Total Sugars (as invert), min	24.00%

INGREDIENTS

Cane Molasses, Soybean Meal, Monocalcium Phosphate, Magnesium Oxide, Vegetable Oil, Calcium Carbonate, Potassium Chloride, Choline Chloride, Niacin Supplement, Vitamin B12 Supplement, D-Calcium Pantothenate, Vitamin E Supplement, Riboflavin Supplement, Vitamin A Supplement, Menadione Sodium Bisulfite Complex, Vitamin D3 Supplement, Thiamine Mononitrate, Biotin, Ascorbic Acid, Folic Acid, Zinc Hydroxychloride, Manganese Hydroxychloride, Cobalt Glucoheptonate, L-Lysine, Selenium Yeast, Basic Copper Chloride, Lactic Acid, Sodium Selenite, Vitamin C Supplement and Ethylenediamine Dihydriodide

FEEDING RECOMMENDATIONS

Consumption will vary depending upon the condition of the animals, quality and quantity of forages being fed, weather conditions, and most importantly the feeding location of the tubs with respect to loafing, grazing, feeding, and watering areas.

Tubs should be distributed at the rate of one tub per 2 pasture horses. Place the tubs where animals congregate (loafing, grazing, feeding, and watering areas). Feed the tubs continuously along with plentiful sources of forage and clean, fresh water. Provide free-choice salt at all times. When feedstuffs are limited, additional nutrients may be required. Contact your feed representative for specific recommendations.

INTRODUCTION PERIOD: During the first week of use, place the tubs approximately 100 feet from loafing and/or watering areas. As livestock become acclimated to the presence of the tubs, repositioning may be necessary to achieve proper consumption. Move tubs closer to increase, further away to decrease consumption. If overconsumption occurs for more than two weeks and repositioning does not correct the situation, remove tubs and evaluate the overall feeding program.

NO SULFATES

NO OXIDES

Manufactured By:
Hudson Livestock Supplements, Inc.
8150 Thompson Rd
Miles, TX 76861
(325) 659-3992 * 1-800-750-9608

NET WEIGHT 125 lb (56.70 kg)