



STUBBLE BUSTER

28-7



FOR BEEF CATTLE ON NATIVE AND IMPROVED PASTURE

GUARANTEED ANALYSIS

Crude Protein, min	28.00%	Potassium (K), min	1.80%
Not more than 7% equivalent protein from non-protein nitrogen		Zinc (Z), min	400 ppm
Crude Fat, min.....	4.00%	Manganese (Mn), min	320 ppm
Crude Fiber, max.....	2.50%	Copper (Cu), min	100 ppm
Calcium (Ca), min	2.00%	Cobalt (Co), min	1 ppm
Calcium (Ca), max	3.00%	Iodine (I), min	7 ppm
Phosphorus (P), min	1.00%	Selenium (Se), min.....	5 ppm
Salt (NaCl)	0 added	Vitamin A, min	100,000 IU/lb
Magnesium (Mg), min.....	1.00%	Vitamin D, min	10,000 IU/lb
		Vitamin E, min.....	20 IU/lb
		Total Sugars (as invert), min	25.00%

INGREDIENTS

Cane Molasses, Hydrolyzed Feather Meal, Cottonseed Meal, Monocalcium Phosphate, Vegetable Oil, Urea, Calcium Carbonate, Magnesium Oxide, Vitamin A Supplement, Vitamin D3 Supplement, Vitamin E Supplement, Selenium Yeast, Manganese Hydroxychloride, Zinc Hydroxychloride, Basic Copper Chloride, Sodium Selenite, Propionic Acid (preservative), Cobalt Carbonate and Ethylenediamine Dihydriodide

Manufactured By:
Hudson Livestock Supplements, Inc.
8150 Thompson Rd
Miles, TX 76861
(325) 659-3992 * 1-800-750-9608

FEEDING RECOMMENDATIONS

Consumption will vary depending upon the condition of the animals, quality and quantity of forages being fed, weather conditions, and most importantly the feeding location of the tubs with respect to loafing, grazing, feeding, and watering areas.

Cattle will consume approximately 1/2 to 1 pound per head daily based upon a 1,000 lb. animal. Tubs should be distributed at the rate of one tub per 20-25 head. Place the tubs where animals congregate (loafing, grazing, feeding, and watering areas). Feed the tubs continuously along with plentiful sources of forage and clean, fresh water. Provide free-choice salt at all times. When feedstuffs are limited, additional nutrients may be required. Contact your feed representative for specific recommendations.

INTRODUCTION PERIOD: During the first week of use, place the tubs approximately 100 feet from loafing and/or watering areas. As livestock become acclimated to the presence of the tubs, repositioning may be necessary to achieve proper consumption. Move tubs closer to increase, further away to decrease consumption. If overconsumption occurs for more than two weeks and repositioning does not correct the situation, remove tubs and evaluate the overall feeding program.

CAUTION:

Use as directed. For ruminants only. Do not feed to equine. Do not feed to sheep or any species that have a low tolerance to supplemental copper.

NO SULFATES

NO OXIDES

NET WEIGHT 225 lb (102.06 kg)